



Chronic Kidney Disease (CKD) Is a Common Comorbidity in Patients With Type 2 Diabetes (T2D)



CKD in T2D is driven by the combined effects of 3 major factors^{4,5}



METABOLIC FACTORS^{4,6}

Controlled by glucose-lowering agents



HEMODYNAMIC FACTORS^{4,6}

Managed by blood pressure lowering agents



INFLAMMATORY AND FIBROTIC FACTORS^{4,5}

Largely unaddressed in patients with CKD in T2D



*From a prospective cohort study of 543,412 adults in Taiwan between 1994 and 2008.

1. American Diabetes Association. Statistics about diabetes. <https://www.diabetes.org/resources/statistics/statistics-about-diabetes>. Accessed May 27, 2020.

2. US Department of Health and Human Services, Centers for Disease Control and Prevention. National diabetes statistics report, 2020.

3. Wen CP, Chang CH, Tsai MK, et al. *Kidney Int.* 2017;92(2):388-396. 4. Alicic RZ, et al. *Clin J Am Soc Nephrol.* 2017;12(12):2032-2045.

5. Bauersachs J, et al. *Hypertension.* 2015;65(2):257-263. 6. American Diabetes Association. *Diabetes Care.* 2020;43(Suppl 1):S135-S151.

To learn more about CKD in T2D visit: www.ckd-t2d.com

