



Chronic Kidney Disease (CKD) Is a Common Comorbidity in Patients With Type 2 Diabetes (T2D)



CKD in T2D is driven by the combined effects of 3 major factors^{4,5}



METABOLIC FACTORS^{4,6}

Controlled by glucose-lowering agents



HEMODYNAMIC FACTORS^{4,6}

Managed by blood pressure-lowering agents



INFLAMMATORY AND FIBROTIC FACTORS^{4,5}

Currently unaddressed in patients with CKD in T2D



^aFrom a prospective cohort study of 543,412 adults in Taiwan between 1994 and 2008.

1. American Diabetes Association. Statistics about diabetes. <https://www.diabetes.org/resources/statistics/statistics-about-diabetes>. Accessed May 27, 2020.

2. US Department of Health and Human Services, Centers for Disease Control and Prevention. National diabetes statistics report, 2020.

3. Wen CP, et al. *Kidney Int.* 2017;92(2):388-396. 4. Alicic RZ, et al. *Clin J Am Soc Nephrol.* 2017;12(12):2032-2045.

5. Bauersachs J, et al. *Hypertension.* 2015;65(2):257-263. 6. American Diabetes Association. *Diabetes Care.* 2020;43(Suppl 1):S135-S151.

To learn more about CKD in T2D visit: www.ckd-t2d.com

